

# **Product datasheet for TA381400**

## **SET Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200 IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 146-277 of

human SET/TAF1 (NP\_003002.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size:30kDa/31kDa/32kDa/33kDaGene Name:SET nuclear proto-oncogene

Database Link: Entrez Gene 6418 Human

Q01105



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

The protein encoded by this gene inhibits acetylation of nucleosomes, especially histone H4, by histone acetylases (HAT). This inhibition is most likely accomplished by masking histone lysines from being acetylated, and the consequence is to silence HAT-dependent transcription. The encoded protein is part of a complex localized to the endoplasmic reticulum but is found in the nucleus and inhibits apoptosis following attack by cytotoxic T lymphocytes. This protein can also enhance DNA replication of the adenovirus genome. Several transcript variants encoding different isoforms have been found for this gene.

Synonyms:

2PP2A; I-2PP2A; I2PP2A; IGAAD; IPP2A2; PHAPII; TAF-I; TAF-IBETA

## **Product images:**



Western blot analysis of extracts of various cell lines, using SET/TAF1 antibody (TA381400) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.