

# Product datasheet for TA381490S

## SIGLECL1 (SIGLEC12) Rabbit Polyclonal Antibody

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Mouse
Modifications:	Unmodified
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 20-210 of human SIGLEC12 (NP_443729.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	51kDa/64kDa
Gene Name:	sialic acid binding lg like lectin 12 (gene/pseudogene)
Database Link:	<u>Q96PQ1</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **GRIGENE** SIGLECL1 (SIGLEC12) Rabbit Polyclonal Antibody – TA381490S

Background:Sialic acid-binding immunoglobulin-like lectins (SIGLECs) are a family of cell surface proteins<br/>belonging to the immunoglobulin superfamily. They mediate protein-carbohydrate<br/>interactions by selectively binding to different sialic acid moieties present on glycolipids and<br/>glycoproteins. This gene encodes a member of the SIGLEC3-like subfamily of SIGLECs.<br/>Members of this subfamily are characterized by an extracellular V-set immunoglobulin-like<br/>domain followed by two C2-set immunoglobulin-like domains, and the cytoplasmic tyrosine-<br/>based motifs ITIM and SLAM-like. The encoded protein, upon tyrosine phosphorylation, has<br/>been shown to recruit the Src homology 2 domain-containing protein-tyrosine phosphatases<br/>SHP1 and SHP2. It has been suggested that the protein is involved in the negative regulation<br/>of macrophage signaling by functioning as an inhibitory receptor. This gene is located in a<br/>cluster with other SIGLEC3-like genes on 19q13.4. Alternative splicing results in multiple<br/>transcript variants.

Synonyms: FLJ38600; S2V; Siglec-12; Siglec-L1; Siglec-XII; SIGLECL1; SLG

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US