

Product datasheet for TA381527S

SKP2 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	ELISA, ICC/IF, WB
Recommended Dilution:	WB,1:100 - 1:500 IF/ICC,1:50 - 1:200 ELISA,Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Reactivity:	Human
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	48kDa
Gene Name:	S-phase kinase-associated protein 2, E3 ubiquitin protein ligase
Database Link:	<u>Entrez Gene 6502 Human</u> <u>Q13309</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE SKP2 Rabbit Polyclonal Antibody – TA381527S

Background:This gene encodes a member of the F-box protein family which is characterized by an
approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four
subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in
phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes:
Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing
either different protein-protein interaction modules or no recognizable motifs. The protein
encoded by this gene belongs to the Fbls class; in addition to an F-box, this protein contains
10 tandem leucine-rich repeats. This protein is an essential element of the cyclin A-CDK2 S-
phase kinase. It specifically recognizes phosphorylated cyclin-dependent kinase inhibitor 1B
(CDKN1B, also referred to as p27 or KIP1) predominantly in S phase and interacts with S-
phase kinase-associated protein 1 (SKP1 or p19). In addition, this gene is established as a
protooncogene causally involved in the pathogenesis of lymphomas. Alternative splicing of
this gene generates three transcript variants encoding different isoforms.

Synonyms:

FBL1; FBXL1; FLB1; MGC1366; p45skp2

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US