

Product datasheet for TA381537S

i roddet datasneet for 1A301337.

SLC10A2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Reactivity: WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 289-348 of

human SLC10A2 (NP_000443.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 37kDa

Gene Name: solute carrier family 10 member 2

Database Link: Entrez Gene 6555 Human

Q12908

Background: This gene encodes a sodium/bile acid cotransporter. This transporter is the primary

mechanism for uptake of intestinal bile acids by apical cells in the distal ileum. Bile acids are the catabolic product of cholesterol metabolism, so this protein is also critical for cholesterol

homeostasis. Mutations in this gene cause primary bile acid malabsorption (PBAM);

muatations in this gene may also be associated with other diseases of the liver and intestines,

such as familial hypertriglyceridemia (FHTG).

Synonyms: ASBT; IBAT; ISBT; NTCP2




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of extracts of various cell lines, using SLC10A2 antibody ([TA381537]) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.