

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA381570

## SLC1A5 Rabbit Polyclonal Antibody


#### **Product data:**

| Product Type:           | Primary Antibodies   |
|-------------------------|--|
| Applications:           | WB   |
| Recommended Dilution:   | WB,1:500 - 1:3000  |
| Reactivity:             | Human, Mouse, Rat  |
| Modifications:          | Unmodified   |
| Host:                   | Rabbit   |
| lsotype:                | IgG  |
| Clonality:              | Polyclonal   |
| Immunogen:              | Recombinant fusion protein containing a sequence corresponding to amino acids 154-224 of human SLC1A5 (NP_005619.1).   |
| Formulation:            | Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.  |
| Concentration:          | lot specific   |
| Purification:           | Affinity purification  |
| Conjugation:            | Unconjugated   |
| Storage:                | Store at -20°C. Avoid freeze / thaw cycles.  |
| Stability:              | Shelf life: one year from despatch.  |
| Predicted Protein Size: | 33kDa/36kDa/56kDa  |
| Gene Name:              | solute carrier family 1 member 5   |
| Database Link:          | <u>Entrez Gene 6510 Human</u><br><u>Q15758</u>   |
| Background:             | The SLC1A5 gene encodes a sodium-dependent neutral amino acid transporter that can act<br>as a receptor for RD114/type D retrovirus (Larriba et al., 2001 [PubMed 11781704]).[supplied<br>by OMIM, Jan 2011] |
| Synonyms:               | AAAT; ASCT2; ATB(0); ATBO; FLJ31068; M7V1; M7VS1; R16; RDR; RDRC   |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **Product images:**



Western blot analysis of extracts of various cell lines, using SLC1A5 antibody (TA381570) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 180s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US