

Product datasheet for TA381874

rioduct datasileet for TASO107.

SOCS4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human, Mouse

Modifications: Unmodified

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-270 of

human SOCS4 (NP_543143.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 50kDa

Gene Name: suppressor of cytokine signaling 4

Database Link: Entrez Gene 122809 Human

Q8WXH5

Background: The protein encoded by this gene contains a SH2 domain and a SOCS BOX domain. The

protein thus belongs to the suppressor of cytokine signaling (SOCS), also known as STAT-induced STAT inhibitor (SSI), protein family. SOCS family members are known to be cytokine-inducible negative regulators of cytokine signaling. Two alternatively spliced transcript

variants encoding the same protein have been found for this gene.

Synonyms: CIS4; DKFZp686J1568; SOCS-4; SOCS-7; SOCS7; SSI4; STAT4




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of extracts of various cell lines, using SOCS4 antibody (TA381874) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.