

## **Product datasheet for TA381905**

## SP110 Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WE

**Reactivity:** WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 1-290 of

human SP110 (NP\_004501.3).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 28kDa/46kDa/61kDa/62kDa/78kDa/81kDa

Gene Name: SP110 nuclear body protein

Database Link: Entrez Gene 3431 Human

Q9HB58

**Background:** The nuclear body is a multiprotein complex that may have a role in the regulation of gene

transcription. This gene is a member of the SP100/SP140 family of nuclear body proteins and

encodes a leukocyte-specific nuclear body component. The protein can function as an

activator of gene transcription and may serve as a nuclear hormone receptor coactivator. In addition, it has been suggested that the protein may play a role in ribosome biogenesis and in the induction of myeloid cell differentiation. Alternative splicing has been observed for this

gene and three transcript variants, encoding distinct isoforms, have been identified.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: FLJ22835; IFI41; IFI75; IPR1; VODI

## **Product images:**



Western blot analysis of extracts of various cell lines, using SP110 antibody (TA381905) at 1:1000 dilution.\_Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.\_Lysates/proteins: 25ug per lane.\_Blocking buffer: 3% nonfat dry milk in TBST.\_Detection: ECL Enhanced Kit .\_Exposure time: 30s.