

Product datasheet for TA381985S

SRC Rabbit Polyclonal Antibody

Product data:

Isotype:

Product Type: Primary Antibodies

ICC/IF, IHC, WB **Applications:**

Recommended Dilution: WB.1:500 - 1:2000

IHC,1:50 - 1:200 IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit **IgG**

Clonality: Polyclonal

Recombinant fusion protein containing a sequence corresponding to amino acids 1-80 of Immunogen:

human Src (NP_005408.1).

Formulation: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 59kDa/60kDa

Gene Name: SRC proto-oncogene, non-receptor tyrosine kinase

Database Link: Entrez Gene 6714 Human

P12931

Background: This gene is highly similar to the v-src gene of Rous sarcoma virus. This proto-oncogene may

> play a role in the regulation of embryonic development and cell growth. The protein encoded by this gene is a tyrosine-protein kinase whose activity can be inhibited by phosphorylation by c-SRC kinase. Mutations in this gene could be involved in the malignant progression of colon cancer. Two transcript variants encoding the same protein have been found for this

gene.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

ASV; c-src; OTTHUMP00000174476; OTTHUMP00000174477; p60-Src; pp60c-src; SRC1

Product images:




Western blot analysis of extracts of various cell lines, using Src antibody ([TA381985]) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 5s.



Immunofluorescence analysis of C6 cells using Src antibody ([TA381985]) at dilution of 1:100. Blue: DAPI for nuclear staining.





Immunofluorescence analysis of NIH/3T3 cells using Src antibody ([TA381985]) at dilution of 1:100. Blue: DAPI for nuclear staining.