

## **Product datasheet for TA382100S**

## **STAT5A Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:100 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Phospho Y694

**Host:** Rabbit

Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** A phospho specific peptide corresponding to residues surrounding Y694 of human STAT5.

**Formulation:** PBS with 0.05% proclin300,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 90kDa

**Gene Name:** signal transducer and activator of transcription 5A

Database Link: Entrez Gene 6776 Human

P42229

**Background:** Carries out a dual signal transduction and activation of transcription. Mediates cellular

responses to the cytokine KITLG/SCF and other growth factors. Mediates cellular responses to ERBB4. May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4. Binds to the GAS element and activates PRL-induced transcription. Regulates the expression of milk

proteins during lactation.

**Synonyms:** MGF; STAT5




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**





Western blot analysis of extracts of K562 cells, using Phospho-STAT5-Y694 pAb ([TA382100]) at 1:1000 dilution.K-562 cells were treated by ATP(5 mM) at 30°C for 1 hour. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA. | Detection: ECL Basic Kit. | Exposure time: 1s.

Western blot analysis of extracts of A-431 cells, using Phospho-STAT5-Y694 pAb ([TA382100]) at 1:1000 dilution or STAT5B antibody (A12356).A-431 cells were treated by EGF (100 ng/mL) at 37°C for 30 minutes after serum-starvation overnight. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA. | Detection: ECL Basic Kit. | Exposure time: 60s.