

Product datasheet for TA382246S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Synaptopodin 2 (SYNPO2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:1000

Reactivity: Mouse

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of

human SYNPO2 (NP_597734.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 74kDa/117kDa/119kDa/133kDa/136kDa

Gene Name: synaptopodin 2

Database Link: Q9UMS6

Background: Has an actin-binding and actin-bundling activity. Can induce the formation of F-actin

networks in an isoform-specific manner. At the sarcomeric Z lines is proposed to act as adapter protein that links nascent myofibers to the sarcolemma via ZYX and may play a role in early assembly and stabilization of the Z lines. Involved in autophagosome formation. May play a role in chaperone-assisted selective autophagy (CASA involved in Z lines maintenance in striated muscle under mechanical tension; may link the client-processing CASA chaperone

machinery to a membrane-tethering and fusion complex providing autophagosome membranes (By similarity. Involved in regulation of cell migration. May be a tumor

suppressor.






Synonyms:

DKFZp686G051; Genethonin-2; myopodin

Product images:



Western blot analysis of extracts of mouse skeletal muscle, using SYNPO2 antibody ([TA382246]) at 1:3000 dilution.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Enhanced Kit.|Exposure time: 90s.