

Product datasheet for TA382348

Trouder addition of the second

Product data:

Product Type: Primary Antibodies

TBPL1 Rabbit Polyclonal Antibody

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse

Modifications: Unmodified

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-186 of

human TBPL1 (NP_004856.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 20kDa

Gene Name: TATA-box binding protein like 1

Database Link: Entrez Gene 9519 Human

P62380

Background: This gene encodes a member of the TATA box-binding protein family. TATA box-binding

proteins play a critical role in transcription by RNA polymerase II as components of the transcription factor IID (TFIID) complex. The encoded protein does not bind to the TATA box and initiates transcription from TATA-less promoters. This gene plays a critical role in

spermatogenesis, and single nucleotide polymorphisms in this gene may be associated with

male infertility. Alternatively spliced transcript variants have been observed for this gene, and

a pseudogene of this gene is located on the long arm of chromosome 3.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

MGC:8389; MGC:9620; STUD; TLF; TLP; TRF2

Product images:



Western blot analysis of extracts of 293T cells, using TBPL1 antibody (TA382348) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit. |Exposure time: 30s.



Immunofluorescence analysis of A549 cells using TBPL1 antibody (TA382348). Blue: DAPI for nuclear staining.