

Product datasheet for TA382363S

Telethonin (TCAP) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies Applications: ELISA, WB Recommended Dilution: WB.1:500 - 1:1000 ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. **Reactivity:** Human Modifications: Unmodified Host: Rabbit lgG Isotype: **Clonality:** Polyclonal Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. **Concentration:** lot specific Purification: Affinity purification **Conjugation:** Unconjugated Store at -20°C. Avoid freeze / thaw cycles. Storage: Stability: Shelf life: one year from despatch. **Predicted Protein Size:** 19kDa Gene Name: titin-cap Database Link: Entrez Gene 8557 Human 015273 Background: Sarcomere assembly is regulated by the muscle protein titin. Titin is a giant elastic protein with kinase activity that extends half the length of a sarcomere. It serves as a scaffold to which myofibrils and other muscle related proteins are attached. This gene encodes a protein found in striated and cardiac muscle that binds to the titin Z1-Z2 domains and is a substrate of titin kinase, interactions thought to be critical to sarcomere assembly. Mutations in this gene are associated with limb-girdle muscular dystrophy type 2G. Synonyms: CMD1N; LGMD2G; T-cap; TELE; telethonin



View online »


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



Product images:



Western blot analysis of lysates from wild type (WT) and 293T cells transfected with TCAP using TCAP Rabbit pAb ([TA382363]) at 1:600 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US