

Product datasheet for TA382394

Product datasireet for TA362394

TDP1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-140 of

human TDP1 (NP_060789.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 41kDa/68kDa

Gene Name: tyrosyl-DNA phosphodiesterase 1

Database Link: Entrez Gene 55775 Human

Q9NUW8

Background: The protein encoded by this gene is involved in repairing stalled topoisomerase I-DNA

complexes by catalyzing the hydrolysis of the phosphodiester bond between the tyrosine residue of topoisomerase I and the 3-prime phosphate of DNA. This protein may also remove glycolate from single-stranded DNA containing 3-prime phosphoglycolate, suggesting a role in repair of free-radical mediated DNA double-strand breaks. This gene is a member of the phospholipase D family and contains two PLD phosphodiesterase domains. Mutations in this gene are associated with the disease spinocerebellar ataxia with axonal neuropathy (SCAN1).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

2810481F14Rik; 4921509N21Rik; Al838772; AW493413; E430034L06Rik; MGC112732; MLZ-501; SCAN1

Product images:



lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit .|Exposure time: 90s.



Immunofluorescence analysis of U-2 OS cells using TDP1 Rabbit pAb (TA382394) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Western blot analysis of extracts of various cell lines, using TDP1 Antibody (TA382394) at 1:1000

dilution. | Secondary antibody: HRP Goat Anti-

Rabbit IgG (H+L) at 1:10000

dilution. | Lysates/proteins: 25ug per