

## Product datasheet for **TA382401**

### TEF1 (TEAD1) Rabbit Polyclonal Antibody

#### Product data:


Product Type:	Primary Antibodies
Applications:	ChIP, ICC/IF, IHC, WB
Recommended Dilution:	WB,1:500 - 1:2000 IF,1:50 - 1:200 ChIP,1:50 - 1:200
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 135-215 of human TEAD1 (NP_068780.2).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	40kDa/47kDa
Gene Name:	TEA domain transcription factor 1
Database Link:	<a href="#">Entrez Gene 7003 Human P28347</a>
Background:	This gene encodes a ubiquitous transcriptional enhancer factor that is a member of the TEA/ATTS domain family. This protein directs the transactivation of a wide variety of genes and, in placental cells, also acts as a transcriptional repressor. Mutations in this gene cause Sveinsson's chorioretinal atrophy. Additional transcript variants have been described but their full-length natures have not been experimentally verified.




[View online »](#)

Synonyms: AA; NTEF-1; REF1; TCF-13; TCF13; TEAD-1; TEF-1


### Product images:



Western blot analysis of extracts of various cell lines, using TEAD1 antibody (TA382401) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 120s.



Immunofluorescence analysis of MCF7 cells using TEAD1 antibody (TA382401).



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using TEAD1 antibody (TA382401) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.