

Product datasheet for TA382402

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TEA domain family member 2 (TEAD2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human TEAD2

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 35kDa/49kDa

Gene Name: TEA domain transcription factor 2

Database Link: Entrez Gene 8463 Human

Q15562

Background: Transcription factor which plays a key role in the Hippo signaling pathway, a pathway

involved in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein MST1/MST2, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Acts by mediating gene expression of YAP1 and WWTR1/TAZ, thereby regulating cell proliferation, migration and epithelial mesenchymal transition (EMT induction. Binds to the SPH and GT-IIC 'enhansons' (5'-GTGGAATGT-3'. May be

involved in the gene regulation of neural development. Binds to the M-CAT motif.






Synonyms:

Etdf; ETF; TEAD-2; TEF-4; TEF4

Product images:



Western blot analysis of extracts of A-431 cells, using HSPB7 antibody (TA382402) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit. |Exposure time: 180s.