

# Product datasheet for TA382665

Recommended Dilution: WB,1:500 - 1:2000

## **RANKL (TNFSF11) Rabbit Polyclonal Antibody**

IHC, WB

Rabbit

Polyclonal

lot specific

014788

Affinity purification

27kDa/30kDa/35kDa

Entrez Gene 8600 Human

Unconjugated

lgG

**Primary Antibodies** 

IHC,1:50 - 1:200 IF,1:50 - 1:200

Human, Mouse

Recombinant protein of human TNFSF11.

Store at -20°C. Avoid freeze / thaw cycles.

tumor necrosis factor superfamily member 11

Shelf life: one year from despatch.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Product data:**

**Product Type:** 

**Applications:** 

**Reactivity:** 

Host:

Isotype:

**Clonality:** 

Immunogen:

Formulation:

Purification:

**Conjugation:** 

Gene Name:

Database Link:

**Predicted Protein Size:** 

Storage: Stability:

Concentration:

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **GRIGENE** RANKL (TNFSF11) Rabbit Polyclonal Antibody – TA382665

**Background:** This gene encodes a member of the tumor necrosis factor (TNF) cytokine family which is a ligand for osteoprotegerin and functions as a key factor for osteoclast differentiation and activation. This protein was shown to be a dentritic cell survival factor and is involved in the regulation of T cell-dependent immune response. T cell activation was reported to induce expression of this gene and lead to an increase of osteoclastogenesis and bone loss. This protein was shown to activate antiapoptotic kinase AKT/PKB through a signaling complex involving SRC kinase and tumor necrosis factor receptor-associated factor (TRAF) 6, which indicated this protein may have a role in the regulation of cell apoptosis. Targeted disruption of the related gene in mice led to severe osteopetrosis and a lack of osteoclasts. The deficient mice exhibited defects in early differentiation of T and B lymphocytes, and failed to form lobulo-alveolar mammary structures during pregnancy. Two alternatively spliced transcript variants have been found.

Synonyms:

CD254; hRANKL2; ODF; OPGL; OPTB2; OTTHUMP00000178585; RANKL; sOdf; TRANCE

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US