

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA382679S

Cardiac Troponin I (TNNI3) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ICC/IF, IP, WB
Recommended Dilution:	WB,1:500 - 1:1000 IF/ICC,1:50 - 1:200 IP,0.5µg-4µg antibody for 400µg-600µg extracts of whole cells ELISA,Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Reactivity:	Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	24kDa
Gene Name:	troponin I3, cardiac type
Database Link:	<u>P19429</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


Cardiac Troponin I (TNNI3) Rabbit Polyclonal Antibody – TA382679S

Background:Troponin I (TnI), along with troponin T (TnT) and troponin C (TnC), is one of 3 subunits that
form the troponin complex of the thin filaments of striated muscle. TnI is the inhibitory
subunit; blocking actin-myosin interactions and thereby mediating striated muscle relaxation.
The TnI subfamily contains three genes: TnI-skeletal-fast-twitch, TnI-skeletal-slow-twitch, and
TnI-cardiac. This gene encodes the TnI-cardiac protein and is exclusively expressed in cardiac
muscle tissues. Mutations in this gene cause familial hypertrophic cardiomyopathy type 7
(CMH7) and familial restrictive cardiomyopathy (RCM). Troponin I is useful in making a
diagnosis of heart failure, and of ischemic heart disease. An elevated level of troponin is also
now used as indicator of acute myocardial injury in patients hospitalized with
moderate/severe Coronavirus Disease 2019 (COVID-19). Such elevation has also been
associated with higher risk of mortality in cardiovascular disease patients hospitalized due to
COVID-19.

Synonyms:


CMD2A; CMH7; cTnl; MGC116817; RCM1; TNNC1

Product images:




Western blot analysis of lysates from Mouse heart

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Western blot analysis of lysates from Rat lung

Immunoprecipitation analysis of 600 µg extracts of Mouse heart cells using 3 µg Cardiac Troponin I (TNNI3) antibody ([TA382679]). Western blot was performed from the immunoprecipitate using Cardiac Troponin I (TNNI3) antibody ([TA382679]) at a dilution of 1:1000.



Immunofluorescence analysis of paraffinembedded mouse heart using Cardiac Troponin I (TNNI3) Rabbit pAb ([TA382679]) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US