

Product datasheet for **TA382721**

p53 (TP53) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IP, WB
Recommended Dilution:	WB,1:500 - 1:2000 IP,1:50 - 1:100
Reactivity:	Human
Modifications:	Phospho S9
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	A synthetic phosphorylated peptide around S9 of human p53 (NP_000537.3).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	23-43kDa
Gene Name:	tumor protein p53
Database Link:	Entrez Gene 7157 Human P04637



[View online »](#)


Background:

This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons from identical transcript variants (PMIDs: 12032546, 20937277).


Synonyms:

FLJ92943; LFS1; p53; TRP53

Product images:



Western blot analysis of extracts of 293 and HT-29 cells, using Phospho-p53-S9 antibody (TA382721) at 1:1000 dilution. 293 cells were treated by UV for 15-30 minutes. HT-29 cells were treated by UV for 15-30 minutes. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% BSA. |Detection: ECL Basic Kit . |Exposure time: 3min.



Immunoprecipitation analysis of 200ug extracts of 293T cells, using 3 ug Phospho-p53-S9 pAb (TA382721). Western blot was performed from the immunoprecipitate using Phospho-p53-S9 pAb (TA382721) at a dilution of 1:1000. 293T cells were treated by UV at room temperature for 30 minutes after serum-starvation overnight, and then treated by 10% FBS at 37°C for 30 minutes.