

Product datasheet for TA382726S

p53 (TP53) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human

Modifications: Phospho S6

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A phospho specific peptide corresponding to residues surrounding S6 of human p53.

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 43kDa

Gene Name: tumor protein p53

Database Link: Entrez Gene 7157 Human

P04637

Background: This gene encodes a tumor suppressor protein containing transcriptional activation, DNA

binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome.

Alternative splicing of this gene and the use of alternate promoters result in multiple

transcript variants and isoforms. Additional isoforms have also been shown to result from the

use of alternate translation initiation codons from identical transcript variants (PMIDs:

12032546, 20937277).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

FLJ92943; LFS1; p53; TRP53

Product images:



Western blot analysis of extracts of MCF7 cells, using Phospho-Tp53-S6 antibody ([TA382726]) at 1:1000 dilution.MCF7 cells were treated by ATP(5 mM) at 30°C for 1 hour. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% BSA. |Detection: ECL Basic Kit. |Exposure time: 90s.