

## **Product datasheet for TA382757S**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

## **Tropomyosin 3 (TPM3) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

**Reactivity:** Mouse

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-284 of

human TPM3 (NP\_689476.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 18kDa/28kDa/29kDa/32kDa

**Gene Name:** tropomyosin 3

Database Link: P06753

**Background:** This gene encodes a member of the tropomyosin family of actin-binding proteins.

Tropomyosins are dimers of coiled-coil proteins that provide stability to actin filaments and regulate access of other actin-binding proteins. Mutations in this gene result in autosomal

dominant nemaline myopathy and other muscle disorders. This locus is involved in

translocations with other loci, including anaplastic lymphoma receptor tyrosine kinase (ALK) and neurotrophic tyrosine kinase receptor type 1 (NTRK1), which result in the formation of fusion proteins that act as oncogenes. There are numerous pseudogenes for this gene on

different chromosomes. Alternative splicing results in multiple transcript variants.






Synonyms:

FLJ41118; Gamma-tropomyosin; hscp30; hTM5; MGC3261; MGC14582; MGC72094; NEM1; OK/SW-cl.5; TM-5; TM3; TM30; TM30nm; TPMsk3; TRK; Tropomyosin-3; Tropomyosin-5

## **Product images:**



Western blot analysis of extracts of various cell lines, using TPM3 antibody ([TA382757]) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST.