

## **Product datasheet for TA382822S**

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **TRIM37 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, WB

Recommended Dilution: WB,1:500 - 1:2000

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

**Reactivity:** Mouse, Rat

**Host:** Rabbit

**Isotype:** IgG

**Clonality:** Polyclonal

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 108kDa

**Gene Name:** tripartite motif containing 37

Database Link: 094972

**Background:** This gene encodes a member of the tripartite motif (TRIM) family, whose members are

involved in diverse cellular functions such as developmental patterning and oncogenesis. The TRIM motif includes zinc-binding domains, a RING finger region, a B-box motif and a coiled-coil domain. The RING finger and B-box domains chelate zinc and might be involved in protein-protein and/or protein-nucleic acid interactions. Mutations in this gene are associated with mulibrey (muscle-liver-brain-eye) nanism, an autosomal recessive disorder that involves several tissues of mesodermal origin. TRIM37 localizes in peroxisomal membranes, and has


been implicated in human peroxisomal biogenesis disorders.

Synonyms: KIAA0898; MUL; POB1; TEF3





## **Product images:**



Western blot analysis of various lysates using TRIM37 Rabbit pAb ([TA382822]) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.