

Product datasheet for TA382974

Tyrosinase (TYR) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IP, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200 IP,1:50 - 1:200

Reactivity: Human, Rat

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human TYR.

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 42kDa/60kDa

Gene Name: tyrosinase

Database Link: Entrez Gene 7299 Human

P14679

Background: The enzyme encoded by this gene catalyzes the first 2 steps, and at least 1 subsequent step,

in the conversion of tyrosine to melanin. The enzyme has both tyrosine hydroxylase and dopa oxidase catalytic activities, and requires copper for function. Mutations in this gene result in oculocutaneous albinism, and nonpathologic polymorphisms result in skin

pigmentation variation. The human genome contains a pseudogene similar to the 3' half of

this gene.

Synonyms: LB24-AB; OCA1A; OCAIA; SHEP3; SK29-AB; tyrosinase




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of extracts of various cell lines, using TYR Rabbit pAb (TA382974) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 60s.