

## **Product datasheet for TA383022S**

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## **UBE2N Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, WB

Recommended Dilution: WB,1:1000 - 1:2000

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 17kDa

**Gene Name:** ubiquitin conjugating enzyme E2 N

Database Link: Entrez Gene 7334 Human

P61088

**Background:** The modification of proteins with ubiquitin is an important cellular mechanism for targeting

abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. Studies in mouse suggest that this protein plays a role in DNA


postreplication repair.

Synonyms: BLU; MGC8489; MGC131857; UBC13; UbcH-ben





## **Product images:**



Western blot analysis of various lysates using Ube2N/Ubc13 Rabbit pAb ([TA383022]) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 15s.