

Product datasheet for TA383046S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Ubiquilin (UBQLN1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB,1:200 - 1:2000

Reactivity: Mouse, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 480-540 of

human UBQLN1 (NP_038466.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 18kDa/43kDa/59kDa/62kDa

Gene Name: ubiquilin 1

Database Link: Q9UMX0

Background: This gene encodes an ubiquitin-like protein (ubiquilin) that shares a high degree of similarity

with related products in yeast, rat and frog. Ubiquilins contain an N-terminal ubiquitin-like domain and a C-terminal ubiquitin-associated domain. They physically associate with both

proteasomes and ubiquitin ligases, and thus are thought to functionally link the

ubiquitination machinery to the proteasome to affect in vivo protein degradation. This ubiquilin has also been shown to modulate accumulation of presenilin proteins, and it is found in lesions associated with Alzheimer's and Parkinson's disease. Two transcript variants

encoding different isoforms have been found for this gene.






Synonyms:

DA41; DSK2; FLJ90054; hPLIC-1; PLIC-1; PLIC1; UBQN; XDRP1

Product images:



Western blot analysis of extracts of various cell lines, using UBQLN1 antibody ([TA383046]) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s