

Product datasheet for TA383048

UBQLN3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 306-620 of

human UBQLN3 (NP_059509.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 70kDa

Gene Name: ubiquilin 3

Database Link: Entrez Gene 50613 Human

Q9H347



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:


This gene encodes a ubiquitin-like protein (ubiquilin) that shares a high degree of similarity with related products in yeast, rat and frog. Ubiquilins contain an N-terminal ubiquitin-like domain and a C-terminal ubiquitin-associated domain. They physically associate with both proteasomes and ubiquitin ligases, and are thus thought to functionally link the ubiquitination machinery to the proteasome to affect in vivo protein degradation. This gene is specifically expressed in the testis. It has been suggested that this gene may regulate cell-cycle progression during spermatogenesis, however, it has been shown that the ortholgous mouse gene is dispensable for embryonic development and spermatogenesis. [provided by RefSeq, Nov 2016]

Synonyms: TUP-1

Product images:




Western blot analysis of extracts of various cell lines, using UBQLN3 antibody (TA383048) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.



Immunofluorescence analysis of rat testis using UBQLN3 antibody (TA383048) at dilution of 1:100. Blue: DAPI for nuclear staining.





Immunofluorescence analysis of mouse testis using UBQLN3 antibody (TA383048) at dilution of 1:100. Blue: DAPI for nuclear staining.