

Product datasheet for **TA383049**

UBQLN4 Rabbit Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	ELISA, ICC/IF, IHC, WB
Recommended Dilution:	WB, 1:500 - 1:1000 IHC-P, 1:50 - 1:200 IF/ICC, 1:50 - 1:200 ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Formulation:	Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	64kDa
Gene Name:	ubiquilin 4
Database Link:	Entrez Gene 56893 Human Q9NRR5
Background:	Enables K48-linked polyubiquitin modification-dependent protein binding activity and identical protein binding activity. Involved in cellular response to DNA damage stimulus; negative regulation of double-strand break repair via homologous recombination; and regulation of cellular catabolic process. Located in several cellular components, including autophagosome; nucleoplasm; and site of DNA damage. Part of protein-containing complex. Colocalizes with cytosolic proteasome complex and nuclear proteasome complex.



[View online »](#)

Synonyms: A1U; C1orf6; CIP75; UBIN

Product images:



Western blot analysis of various lysates using UBQLN4 Rabbit pAb (TA383049) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 60s.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using UBQLN4 Rabbit pAb (TA383049) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human placenta using UBQLN4 Rabbit pAb (TA383049) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse lung using UBQLN4 Rabbit pAb (TA383049) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain using UBQLN4 Rabbit pAb (TA383049) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of PC-12 cells using UBQLN4 Rabbit pAb (TA383049) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.