

Product datasheet for TA383086

Uhrf2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB.1:500 - 1:2000

Reactivity: Mouse, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: **IgG**

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 76-359 of

mouse Uhrf2 (NP_659122.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 24kDa/57kDa/90kDa

Gene Name: ubiquitin-like, containing PHD and RING finger domains 2

Database Link: Entrez Gene 109113 Mouse

Q7TMI3

Background: This gene encodes a nuclear protein which is involved in cell-cycle regulation. The encoded

> protein is a ubiquitin-ligase capable of ubiquinating PCNP (PEST-containing nuclear protein), and together they may play a role in tumorigenesis. The encoded protein contains an NIRF N domain, a PHD finger, a set- and ring-associated (SRA) domain, and a RING finger domain and several of these domains have been shown to be essential for the regulation of cell proliferation. This protein may also have a role in intranuclear degradation of polyglutamine aggregates. Alternative splicing results in multiple transcript variants some of which are non-

protein coding.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

DKFZp434B0920; DKFZp686G0837; MGC33463; NIRF; RNF107; URF2

Product images:



Western blot analysis of extracts of various cell lines, using Uhrf2 antibody (TA383086) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 60s.