

Product datasheet for TA383101S

UMPS Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:100

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 100-260 of

human UMPS (NP_000364.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 22kDa/33kDa/42kDa/52kDa

Gene Name: uridine monophosphate synthetase

Database Link: Entrez Gene 7372 Human

P11172

Background: This gene encodes a uridine 5'-monophosphate synthase. The encoded protein is a

bifunctional enzyme that catalyzes the final two steps of the de novo pyrimidine biosynthetic

pathway. The first reaction is carried out by the N-terminal enzyme orotate

phosphoribosyltransferase which converts orotic acid to orotidine-5'-monophosphate. The terminal reaction is carried out by the C-terminal enzyme OMP decarboxylase which converts orotidine-5'-monophosphate to uridine monophosphate. Defects in this gene are the cause

of hereditary orotic aciduria. Alternate splicing results in multiple transcript variants.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: OMPdecase; OPRT; OPRTase

Product images:



Western blot analysis of extracts of various cell lines, using UMPS antibody ([TA383101]) at 1:3000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit. |Exposure time: 30s.



Immunofluorescence analysis of NIH-3T3 cells using UMPS Polyclonal Antibody ([TA383101]) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.