

## **Product datasheet for TA383113**

## **UPF2 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** IP, WB

Recommended Dilution: WB.1:500 - 1:2000

Reactivity: Human, Mouse

**Modifications:** Unmodified

Host: Rabbit

Isotype: **IgG** 

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1153-1272

of human UPF2 (NP\_542166.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** 66kDa/147kDa

UPF2 regulator of nonsense transcripts homolog (yeast) Gene Name:

**Database Link:** Entrez Gene 26019 Human

Q9HAU5

Background: This gene encodes a protein that is part of a post-splicing multiprotein complex involved in

> both mRNA nuclear export and mRNA surveillance. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD

to degrade mRNAs containing premature stop codons. This protein is located in the perinuclear area. It interacts with translation release factors and the proteins that are

functional homologs of yeast Upf1p and Upf3p. Two splice variants have been found for this

gene; both variants encode the same protein.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

DKFZp434D222; HUPF2; KIAA1408; MGC138834; MGC138835; OTTHUMP00000019091; RENT2; smg-3

## **Product images:**



Western blot analysis of extracts of various cell lines, using UPF2 antibody (TA383113) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.