

Product datasheet for TA383125S

UQCRB Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB.1:1000 - 1:2000 Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: **IgG**

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-111 of

human UQCRB (NP_006285.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 13kDa/16kDa

ubiquinol-cytochrome c reductase binding protein Gene Name:

Database Link: Entrez Gene 7381 Human

P14927

Background: This gene encodes a subunit of the ubiquinol-cytochrome c oxidoreductase complex, which

> consists of one mitochondrial-encoded and 10 nuclear-encoded subunits. The protein encoded by this gene binds ubiquinone and participates in the transfer of electrons when ubiquinone is bound. This protein plays an important role in hypoxia-induced angiogenesis through mitochondrial reactive oxygen species-mediated signaling. Mutations in this gene are associated with mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. Related pseudogenes have been identified on chromosomes

1, 5 and X.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

FLJ92016; FLJ97033; QCR7; QP-C; QPC; UQBC; UQBP; UQPC

Product images:



Western blot analysis of extracts of various cell lines, using UQCRB antibody ([TA383125]) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.