

## **Product datasheet for TA383129S**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **UQCRFS1** Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, IHC, WB

Recommended Dilution: WB,1:500 - 1:1000

IHC-P,1:50 - 1:200

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 30kDa

**Gene Name:** ubiquinol-cytochrome c reductase, Rieske iron-sulfur polypeptide 1

Database Link: Entrez Gene 7386 Human

P47985

**Background:** Predicted to enable oxidoreductase activity. Involved in mitochondrial respiratory chain

complex III assembly and respiratory electron transport chain. Located in mitochondrion. Part of mitochondrial respiratory chain complex III and mitochondrial respiratory chain complex


IV. Implicated in mitochondrial complex III deficiency.


Synonyms: RIP1; RIS1; RISP; UQCR5





## **Product images:**





Western blot analysis of various lysates using UQCRFS1 Rabbit pAb ([TA383129]) at 1:500 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3s.

Immunohistochemistry analysis of paraffinembedded Rat kidney using UQCRFS1 Rabbit pAb ([TA383129]) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.