

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA383131S

URM1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000 IF,1:50 - 1:100
Reactivity:	Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-63 of human URM1 (NP_001252511.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	7kDa/11kDa/15kDa
Gene Name:	ubiquitin related modifier 1
Database Link:	<u>Q9BTM9</u>
Background:	Acts as a sulfur carrier required for 2-thiolation of mcm(5S(2U at tRNA wobble positions of cytosolic tRNA(Lys, tRNA(Glu and tRNA(Gln. Serves as sulfur donor in tRNA 2-thiolation reaction by being thiocarboxylated (-COSH at its C-terminus by MOCS3. The sulfur is then transferred to tRNA to form 2-thiolation of mcm(5S(2U. Also acts as a ubiquitin-like protein (UBL that is covalently conjugated via an isopeptide bond to lysine residues of target proteins such as MOCS3, ATPBD3, CTU2, USP15 and CAS. The thiocarboxylated form serves as substrate for conjugation and oxidative stress specifically induces the formation of UBL-protein conjugates.




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Synonyms:

C9orf74; MGC2668

Product images:



Western blot analysis of extracts of rat eye, using URM1 antibody ([TA383131]) at 1:3000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit . |Exposure time: 90s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US