

### **Product datasheet for TA383383**

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RanBP16 (XPO7) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WE

**Reactivity:** WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthetic peptide corresponding to a sequence within amino acids 1000 to the C-terminus

of human XPO7 (NP\_055839.3).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 123kDa

Gene Name: exportin 7

Database Link: Entrez Gene 23039 Human

Q9UIA9

**Background:** The transport of protein and large RNAs through the nuclear pore complexes (NPC) is an

energy-dependent and regulated process. The import of proteins with a nuclear localization signal (NLS) is accomplished by recognition of one or more clusters of basic amino acids by the importin-alpha/beta complex; see MIM 600685 and MIM 602738. The small GTPase RAN (MIM 601179) plays a key role in NLS-dependent protein import. RAN-binding protein-16 is a


member of the importin-beta superfamily of nuclear transport receptors.

**Synonyms:** Exp7; KIAA0745; RANBP16





# **Product images:**



Western blot analysis of extracts of various cell lines, using XPO7 antibody (TA383383) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.