

## **Product datasheet for TA383399**

## Troduct datasirect for 17,50555

## **YAP1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

**Reactivity:** WB,1:500 - 1:2000 **Reactivity:** Human, Mouse, Rat

Modifications: Phospho S397

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A phospho specific peptide corresponding to residues surrounding S397 of human YAP1.

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 36kDa/48kDa/49kDa/50kDa/52kDa/53kDa/54kDa

Gene Name: Yes associated protein 1

Database Link: Entrez Gene 10413 Human

P46937

**Background:** This gene encodes a downstream nuclear effector of the Hippo signaling pathway which is

involved in development, growth, repair, and homeostasis. This gene is known to play a role in the development and progression of multiple cancers as a transcriptional regulator of this signaling pathway and may function as a potential target for cancer treatment. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by

RefSeq, Aug 2013]

**Synonyms:** YAP; YAP2; YAP65; YKI




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Western blot analysis of extracts of various cell lines, using phospho-YAP1-S397 antibody (TA383399) at 1:1000 dilution.NIH3T3 cells and C6 cells were treated by Sorbitol (0.2M) at 37°C for 60 minutes after serumstarvation overnight. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 1s.