

# **Product datasheet for TA383422**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### 14-3-3 zeta (YWHAZ) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WE

**Reactivity:** WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthetic peptide corresponding to a sequence within amino acids 100 to the C-terminus of

human YWHAZ (NP\_001129171.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 19kDa/27kDa

**Gene Name:** tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta

Database Link: Entrez Gene 7534 Human

P63104

**Background:** This gene product belongs to the 14-3-3 family of proteins which mediate signal transduction

by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals, and this protein is 99% identical to the mouse, rat and sheep orthologs. The encoded protein interacts with IRS1 protein, suggesting a role in regulating insulin sensitivity. Several transcript variants that differ in the 5' UTR but that

encode the same protein have been identified for this gene.






Synonyms:

14-3-3-zeta; KCIP-1; MGC111427; MGC126532; MGC138156; OTTHUMP00000165851; OTTHUMP00000165858

# **Product images:**



Western blot analysis of extracts of various cell lines, using YWHAZ Antibody (TA383422) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.