

Product datasheet for TA383441

ZBTB48 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB.1:500 - 1:2000

Reactivity: Human, Mouse

Modifications: Unmodified

Host: Rabbit

Isotype: **IgG**

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of

human ZBTB48 (NP_005332.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 77kDa

zinc finger and BTB domain containing 48 Gene Name:

Database Link: Entrez Gene 3104 Human

P10074

Background: Telomere-binding protein that acts as a regulator of telomere length. Directly binds the

telomeric double-stranded 5'-TTAGGG-3' repeat. Preferentially binds to telomeres that have a

low concentration of shelterin complex and acts as a regulator of telomere length by initiating telomere trimming, a process that prevents the accumulation of aberrantly long telomeres. Also acts as a transcription regulator that binds to promoter regions. Regulates expression of a small subset of genes, including MTFP1. Regulates expression the J and/or S elements in MHC II promoter. Acts as a negative regulator of cell proliferation by specifically

activating expression of ARF, a tumor suppressor isoform of CDKN2A.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: HKR3; pp9964; ZNF855

Product images:



Western blot analysis of extracts of various cell lines, using ZBTB48 antibody (TA383441) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 90s.