

Product datasheet for TA383509S

PRKCBP1 (ZMYND8) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies Applications: ELISA, WB Recommended Dilution: WB.1:1000 - 1:2000 ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. **Reactivity:** Human Modifications: Unmodified Host: Rabbit lgG Isotype: **Clonality:** Polyclonal Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. **Concentration:** lot specific Purification: Affinity purification **Conjugation:** Unconjugated Store at -20°C. Avoid freeze / thaw cycles. Storage: Shelf life: one year from despatch. Stability: **Predicted Protein Size:** 132kDa Gene Name: zinc finger MYND-type containing 8 Database Link: Entrez Gene 23613 Human Q9ULU4 Background: The protein encoded by this gene is a receptor for activated C-kinase (RACK) protein. The encoded protein has been shown to bind in vitro to activated protein kinase C beta I. In addition, this protein is a cutaneous T-cell lymphoma-associated antigen. Finally, the protein contains a bromodomain and two zinc fingers, and is thought to be a transcriptional regulator. Multiple transcript variants encoding several different isoforms have been found for this gene. Synonyms: KIAA1125; MGC31836; PRKCBP1; PRO2893; RACK7




View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product images:



Western blot analysis of various lysates using ZMYND8 Rabbit pAb ([TA383509]) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 5s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US