

# **Product datasheet for TA383531**

## **ZNF207 Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB,1:1000 - 1:2000

IHC,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant protein of human ZNF207.

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 51kDa

Gene Name: zinc finger protein 207

Database Link: Entrez Gene 7756 Human

<u>043670</u>

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

Kinetochore- and microtubule-binding protein that plays a key role in spindle assembly. ZNF207/BuGZ is mainly composed of disordered low-complexity regions and undergoes phase transition or coacervation to form temperature-dependent liquid droplets. Coacervation promotes microtubule bundling and concentrates tubulin, promoting microtubule polymerization and assembly of spindle and spindle matrix by concentrating its building blocks. Also acts as a regulator of mitotic chromosome alignment by mediating the stability and kinetochore loading of BUB3. Mechanisms by which BUB3 is protected are unclear: according to a first report, ZNF207/BuGZ may act by blocking ubiquitination and proteasomal degradation of BUB3. According to another report, the stabilization is independent of the proteasome.

Synonyms: DKFZp761N202

### **Product images:**



Western blot analysis of extracts of various cell lines, using ZNF207 antibody (TA383531) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: