

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA383565S

ZNF444 Rabbit Polyclonal Antibody

Product data:

| Product Type: | Primary Antibodies |
|-------------------------|--|
| Applications: | ELISA, ICC/IF, IHC, WB |
| Recommended Dilution: | WB,1:500 - 1:2000 IHC-P,1:50 - 1:200 IF/ICC,1:50 - 1:200 ELISA,Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| lsotype: | lgG |
| Clonality: | Polyclonal |
| Formulation: | Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| Concentration: | lot specific |
| Purification: | Affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. |
| Stability: | Shelf life: one year from despatch. |
| Predicted Protein Size: | 35kDa |
| Gene Name: | zinc finger protein 444 |
| Database Link: | <u>Entrez Gene 55311 Human</u> <u>Q8N0Y2</u> |
| Background: | This gene encodes a zinc finger protein which activates transcription of a scavenger receptor gene involved in the degradation of acetylated low density lipoprotein (Ac-LDL) (PMID: 11978792). This gene is located in a cluster of zinc finger genes on chromosome 19 at q13.4. A pseudogene of this gene is located on chromosome 15. Multiple transcript variants encoding different isoforms have been found for this gene. |
| Synonyms: | EZF-2; EZF2; FLJ11137; ZSCAN17 |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Product images:



Western blot analysis of various lysates using ZNF444 Rabbit pAb ([TA383565]) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021). Exposure time: 90s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US