

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA383623

### **ZRANB3 Rabbit Polyclonal Antibody**

## **Product data:**

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:1000 - 1:3000
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 818-1077 of human ZRANB3 (NP_001273497.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	42kDa/49kDa/67kDa/123kDa
Gene Name:	zinc finger RANBP2-type containing 3
Database Link:	<u>Entrez Gene 84083 Human</u> <u>Q5FWF4</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


#### **GRIGENE** ZRANB3 Rabbit Polyclonal Antibody – TA383623

Background:DNA annealing helicase and endonuclease required to maintain genome stability at stalled or<br/>collapsed replication forks by facilitating fork restart and limiting inappropriate<br/>recombination that could occur during template switching events. Recruited to the sites of<br/>stalled DNA replication by polyubiquitinated PCNA and acts as a structure-specific<br/>endonuclease that cleaves the replication fork D-loop intermediate, generating an accessible<br/>3'-OH group in the template of the leading strand, which is amenable to extension by DNA<br/>polymerase. In addition to endonuclease activity, also catalyzes the fork regression via<br/>annealing helicase activity in order to prevent disintegration of the replication fork and the<br/>formation of double-strand breaks.

Synonyms:

4933425L19Rik; DKFZP434B1727; MGC75012; MGC105033

## **Product images:**



Western blot analysis of extracts of various cell lines, using ZRANB3 antibody (TA383623) at 1:3000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit . |Exposure time: 20s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US