

## Product datasheet for TA384065M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Cullin 5 (CUL5) Rabbit Monoclonal Antibody [Clone ID: R02-5K8]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: R02-5K8

Applications: WB

Recommended Dilution: WB: 1/1000-1/5000

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

**Immunogen:** Recombinant protein of human Cullin 5

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

**Concentration:** lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 91 kDa; Observed MW: 91 kDa

**Gene Name:** cullin 5

Database Link: Entrez Gene 8065 Human

Q93034

Background: Swiss-Prot Acc.Q93034.Core component of multiple SCF-like ECS (Elongin-Cullin 2/5-SOCS-box

protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination and

subsequent proteasomal degradation of target proteins. As a scaffold protein may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable substrate recognition component. ECS(SOCS1) seems to direct ubiquitination of JAK2. Seems to be involved in proteosomal degradation of p53/TP53 stimulated by adenovirus E1B-55 kDa

protein. May form a cell surface vasopressin receptor.






Synonyms:

CUL-5; VACM-1; VACM1

# **Product images:**



Western blot analysis of Cullin 5/CUL5 in C6, 3T3, Hela lysates using Cullin 5 antibody.