

### **Product datasheet for TA384078**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cytochrome C (CYCS) Rabbit Monoclonal Antibody [Clone ID: R02-9A7]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: R02-9A7

Applications: IF, IP, WB

Recommended Dilution: WB: 1/1000

ICC/IF: 1/20-1/50

IP: 1/20

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

**Immunogen:** A synthetic peptide of human Cytochrome C

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

**Concentration:** lot specific

Purification:Affinity PurifiedConjugation:Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 12 kDa; Observed MW: 14 kDa

**Gene Name:** cytochrome c, somatic

Database Link: Entrez Gene 54205 Human

P99999






#### Background:

Swiss-Prot Acc.P99999.Electron carrier protein. The oxidized form of the cytochrome c heme group can accept an electron from the heme group of the cytochrome c1 subunit of cytochrome reductase. Cytochrome c then transfers this electron to the cytochrome oxidase complex, the final protein carrier in the mitochondrial electron-transport chain.Plays a role in apoptosis. Suppression of the anti-apoptotic members or activation of the pro-apoptotic members of the Bcl-2 family leads to altered mitochondrial membrane permeability resulting in release of cytochrome c into the cytosol. Binding of cytochrome c to Apaf-1 triggers the activation of caspase-9, which then accelerates apoptosis by activating other caspases.SitesFeature keyPosition(s)Description


Synonyms:

CYC; HCS; OTTHUMP00000202043; OTTHUMP00000202044; THC4

# **Product images:**



Western blot detection of Cytochrome C in K562,Rat Brain,C6,3T3,Hela cell lysates using Cytochrome C Rabbit mAb(1:1000 diluted).Predicted band size:12kDa.Observed band size:14kDa.



Immunocytochemistry analysis of Cytochrome C (green) in K562 using Cytochrome C antibody, and DAPI(blue).