

Product datasheet for TA384080

CYP1B1 Rabbit Monoclonal Antibody [Clone ID: R02-5A3]

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Primary Antibodies |
|-------------------------|--|
| Clone Name: | R02-5A3 |
| Applications: | IHC, WB |
| Recommended Dilution: | WB: 1/2000 IHC: 1/50 |
| Reactivity: | Human, Rat |
| Host: | Rabbit |
| lsotype: | lgG |
| Clonality: | Monoclonal |
| Immunogen: | Recombinant protein of human CYP1B1 |
| Formulation: | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA |
| Concentration: | lot specific |
| Purification: | Affinity Purified |
| Conjugation: | Unconjugated |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Stability: | 1 year |
| Predicted Protein Size: | Calculated MW: 61 kDa; Observed MW: 61 kDa |
| Gene Name: | cytochrome P450 family 1 subfamily B member 1 |
| Database Link: | <u>Entrez Gene 1545 Human</u> <u>Q16678</u> |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE CYP1B1 Rabbit Monoclonal Antibody [Clone ID: R02-5A3] – TA384080

Background:Swiss-Prot Acc.Q16678.Cytochromes P450 are a group of heme-thiolate monooxygenases. In
liver microsomes, this enzyme is involved in an NADPH-dependent electron transport
pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty
acids, retinoid and xenobiotics. Preferentially oxidizes 17beta-estradiol to the carcinogenic 4-
hydroxy derivative, and a variety of procarcinogenic compounds to their activated forms,
including polycyclic aromatic hydrocarbons. Promotes angiogenesis by removing cellular
oxygenation products, thereby decreasing oxidative stress, release of antiangiogenic factor
THBS2, then allowing endothelial cells migration, cell adhesion and capillary morphogenesis.
These changes are concommitant with the endothelial nitric oxide synthase activity and nitric
oxide synthesis. Plays an important role in the regulation of perivascular cell proliferation,
migration, and survival through modulation of the intracellular oxidative state and NF-kappa-
B expression and/or activity, during angiogenesis. Contributes to oxidative homeostasis and
ultrastructural organization and function of trabecular meshwork tissue through modulation
of POSTN expression.

CP1B; CYPIB1; GLC3A; OTTHUMP00000201401; P4501B1


Product images:

Synonyms:



Western blot detection of CYP1B1 in Rat Brain lysates using CYP1B1 Rabbit mAb(1:1000 diluted).Predicted band size:61kDa.Observed band size:61kDa.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of CYP1B1 in paraffinembedded Human breast cancer tissue using CYP1B1 Rabbit mAb at dilution 1/50

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US