

# **Product datasheet for TA384091M**

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## DDX21 Rabbit Monoclonal Antibody [Clone ID: R04-4]2]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: R04-4J2
Applications: IF, WB

Recommended Dilution: WB: 1/1000

ICC/IF: 1/100

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Monoclonal

**Immunogen:** A synthetic peptide of human DDX21

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

**Concentration:** lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

**Storage:** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 87 kDa; Observed MW: 87 kDa

Gene Name: DEAD-box helicase 21

Database Link: Entrez Gene 9188 Human

Q9NR30




#### Background:

Swiss-Prot Acc.Q9NR30.RNA helicase that acts as a sensor of the transcriptional status of both RNA polymerase (Pol) I and II: promotes ribosomal RNA (rRNA) processing and transcription from polymerase II (Pol II) (PubMed:25470060). Binds various RNAs, such as rRNAs, snoRNAs, 7SK and, at lower extent, mRNAs (PubMed:25470060). In the nucleolus, localizes to rDNA locus, where it directly binds rRNAs and snoRNAs, and promotes rRNA transcription, processing and modification. Required for rRNA 2'-O-methylation, possibly by promoting the recruitment of late-acting snoRNAs SNORD56 and SNORD58 with preribosomal complexes (PubMed:25470060, PubMed:25477391). In the nucleoplasm, binds 7SK RNA and is recruited to the promoters of Pol II-transcribed genes: acts by facilitating the release of P-TEFb from inhibitory 7SK snRNP in a manner that is dependent on its helicase activity, thereby promoting transcription of its target genes (PubMed:25470060). Functions as cofactor for JUN-activated transcription: required for phosphorylation of JUN at 'Ser-77' (PubMed:11823437, PubMed:25260534). Can unwind double-stranded RNA (helicase) and can fold or introduce a secondary structure to a single-stranded RNA (foldase) (PubMed:9461305). Involved in rRNA processing (PubMed:14559904, PubMed:18180292). May bind to specific miRNA hairpins (PubMed:28431233).

Synonyms:


DKFZp686F21172; Gu-alpha; GUA; GURDB; RH-II/GU; RH-II/GuA

## **Product images:**



Western blot detection of DDX21 in Hela cell lysates using DDX21 Rabbit mAb(1:1000 diluted). Predicted band size:87kDa. Observed band size:87kDa.





Immunocytochemistry analysis of DDX21 (green) in 293T using DDX21 antibody,and DAPI(blue).