

## Product datasheet for TA384148M

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PERK (EIF2AK3) Rabbit Monoclonal Antibody [Clone ID: R09-1F9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: R09-1F9

Applications: WB

Recommended Dilution: WB: 1/1000

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: Recombinant protein of human PERK

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

**Concentration:** lot specific

Purification: Affinity Purified Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 125 kDa; Observed MW: 140 kDa

**Gene Name:** eukaryotic translation initiation factor 2 alpha kinase 3

Database Link: Entrez Gene 9451 Human

Q9NZJ5






#### Background:

Swiss-Prot Acc.Q9NZJ5.Metabolic-stress sensing protein kinase that phosphorylates the alpha subunit of eukaryotic translation initiation factor 2 (eIF-2-alpha/EIF2S1) on 'Ser-52' during the unfolded protein response (UPR) and in response to low amino acid availability. Converts phosphorylated eIF-2-alpha/EIF2S1 either in a global protein synthesis inhibitor, leading to a reduced overall utilization of amino acids, or to a translation initiation activator of specific mRNAs, such as the transcriptional activator ATF4, and hence allowing ATF4-mediated reprogramming of amino acid biosynthetic gene expression to alleviate nutrient depletion. Serves as a critical effector of unfolded protein response (UPR)-induced G1 growth arrest due to the loss of cyclin-D1 (CCND1). Involved in control of mitochondrial morphology and function.

Synonyms:

DKFZp781H1925; HRI; HsPEK; PEK; PERK; WRS

# **Product images:**



Western blot analysis of PERK in K562, C6, 3T3, Hela lysates using PERK antibody.