

## **Product datasheet for TA384152M**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **EIF3B Rabbit Monoclonal Antibody [Clone ID: R09-2A9]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: R09-2A9
Applications: IHC, WB

Recommended Dilution: WB: 1/1000-1/5000

IHC: 1/20-1/50

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Monoclonal

**Immunogen:** A synthetic peptide of human eIF3B

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

**Concentration:** lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

**Storage:** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 93 kDa; Observed MW: 117 kDa

Gene Name: eukaryotic translation initiation factor 3 subunit B

Database Link: Entrez Gene 8662 Human

P55884




#### Background:

Swiss-Prot Acc.P55884.RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:9388245, PubMed:17581632, PubMed:25849773, PubMed:27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:9388245, PubMed:17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:25849773).

Synonyms:


eIF-3-eta; EIF3-ETA; EIF3-P110; EIF3-P116; EIF3S9; hPrt1; MGC104664; MGC131875; PRT1

#### **Product images:**



Western blot detection of eIF3B in K562,Hela cell lysates using eIF3B Rabbit mAb(1:1000 diluted).Predicted band size:93kDa.Observed band size:117kDa.





Immunohistochemistry of eIF3B in paraffinembedded Human breast cancer tissue using eIF3B Rabbit mAb at dilution 1/20