

Product datasheet for TA384153S

EIF3B Rabbit Monoclonal Antibody [Clone ID: R09-4I5]

Product data:

Product Type: Primary Antibodies Clone Name: R09-415 **Applications:** IF, IP, WB Recommended Dilution: WB: 1/2000-1/10000 ICC/IF: 1/20-1/50 IP: 1/20-1/50 **Reactivity:** Human, Mouse, Monkey Host: Rabbit Isotype: lgG Monoclonal **Clonality:** Immunogen: A synthetic peptide of human eIF3B 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA Formulation: Concentration: lot specific **Purification:** Affinity Purified **Conjugation:** Unconjugated Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Storage: Stability: 1 year Predicted Protein Size: Calculated MW: 93 kDa; Observed MW: 117 kDa Gene Name: eukaryotic translation initiation factor 3 subunit B Database Link: Entrez Gene 8662 Human P55884



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Serigene EIF3B Rabbit Monoclonal Antibody [Clone ID: R09-415] – TA384153S


Background:

Swiss-Prot Acc.P55884.RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:9388245, PubMed:17581632, PubMed:25849773, PubMed:27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:9388245, PubMed:17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:25849773).

Synonyms:

eIF-3-eta; EIF3-ETA; EIF3-P110; EIF3-P116; EIF3S9; hPrt1; MGC104664; MGC131875; PRT1


Product images:



Western blot detection of eIF3B in Raji,COS7,3T3,Hela cell lysates using eIF3B Rabbit mAb(1:1000 diluted).Predicted band size:93KDa.Observed band size:117KDa.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





Immunocytochemistry analysis of eIF3B(green) in Hela using eIF3B antibody,and DAPI(blue)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US