

Product datasheet for TA384173

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Non Neuronal Enolase (ENO1) Rabbit Monoclonal Antibody [Clone ID: R02-2H4]

Product data:

Product Type: Primary Antibodies

Clone Name: R02-2H4

Applications: IF, IP, WB

Recommended Dilution: WB: 1/2000

ICC/IF: 1/50 IP: 1/20

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Monoclonal

Immunogen: A synthetic peptide of human ENO1

Formulation: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

Concentration: lot specific

Purification: Affinity Purified Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Calculated MW: 47 kDa; Observed MW: 47 kDa

Gene Name: enolase 1

Database Link: Entrez Gene 2023 Human

P06733






Background:

Swiss-Prot Acc.P06733.Multifunctional enzyme that, as well as its role in glycolysis, plays a part in various processes such as growth control, hypoxia tolerance and allergic responses. May also function in the intravascular and pericellular fibrinolytic system due to its ability to serve as a receptor and activator of plasminogen on the cell surface of several cell-types such as leukocytes and neurons. Stimulates immunoglobulin production.MBP1 binds to the myc promoter and acts as a transcriptional repressor. May be a tumor suppressor.MiscellaneousUsed as a diagnostic marker for many tumors and, in the heterodimeric form, alpha/gamma, as a marker for hypoxic brain injury after cardiac arrest. Also marker for endometriosis. Antibodies against alpha-enolase are present in sera from patients with cancer-associated retinopathy syndrome (CAR), a progressive blinding disease which occurs in the presence of systemic tumor growth, primarily small-cell carcinoma of the lung and other malignancies. Is identified as an autoantigen in Hashimoto encephalopathy (HE) a rare autoimmune disease associated with Hashimoto thyroiditis (HT). HT is a disorder in which destructive processes overcome the potential capacity of thyroid replacement leading to hypothyroidism.

Synonyms:


ENO1L1; MBP-1; MBPB1; MPB-1; MPB1; NNE; PPH; tau-crystallin

Product images:




Western blot analysis of ENO1 in mouse brain, 3T3, mouse heart lysates using ENO1 antibody.






Western blot analysis of ENO1 in Jurkat lysates using ENO1 antibody.



Western blot analysis of ENO1 in Jurkat, THP-1, C6 lysates using ENO1 antibody



Immunocytochemistry analysis of ENO1(green) in Hela using ENO1 antibody, and DAPI(blue).