

Product datasheet for TA384226M

FGFR1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IF, WB

Recommended Dilution: WB: 1/500-1/2000

IF: 1/200-1/1000 ELISA: 1/20000

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: The antiserum was produced against synthesized peptide derived from human FGFR1

around the phosphorylation site of Tyr654. AA range:626-675 (Phosphorylated)

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Concentration: lot specific

Purification: Affinity Chromatography

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Stability: 1 year

Predicted Protein Size: Observed MW (kDa):120

Database Link: P11362

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


Swiss-Prot Acc.P11362.Tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of embryonic development, cell proliferation, differentiation and migration. Required for normal mesoderm patterning and correct axial organization during embryonic development, normal skeletogenesis and normal development of the gonadotropin-releasing hormone (GnRH) neuronal system. Phosphorylates PLCG1, FRS2, GAB1 and SHB. Ligand binding leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. Phosphorylation of FRS2 triggers recruitment of GRB2, GAB1, PIK3R1 and SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. Promotes phosphorylation of SHC1, STAT1 and PTPN11/SHP2. In the nucleus, enhances RPS6KA1 and CREB1 activity and contributes to the regulation of transcription. FGFR1 signaling is down-regulated by IL17RD/SEF, and by FGFR1 ubiquitination, internalization and degradation.

Product images:




Western blot analysis of Phospho-FGFR1 (Tyr654) in Hela lysates using Phospho-FGFR1 (Tyr654) antibody.







Western blot analysis of Phospho-FGFR1 (Tyr654) in 293 lysates treated with Insulin using Phospho-FGFR1 (Tyr654) antibody. The lane on the right is blocked with the Phospho-peptide.



EnzymeLinked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phospho-peptide (Phospho-left) and NonPhospho-peptide (Phospho-right), using FGFR1 (Phospho-Tyr654) antibody.





Immunofluorescence analysis of Phospho-FGFR1 (Tyr654) in COS7 cells using FGFR1 (Phospho-Tyr654) antibody. The picture on the right is blocked with the Phospho-peptide.