

Product datasheet for **TA384249**

FOXC1 Rabbit Monoclonal Antibody [Clone ID: R06-3B5]

Product data:

Product Type:	Primary Antibodies
Clone Name:	R06-3B5
Applications:	IP, WB
Recommended Dilution:	WB: 1/1000 IP: 1/20
Reactivity:	Mouse
Host:	Rabbit
Isotype:	IgG
Clonality:	Monoclonal
Immunogen:	Recombinant protein of human FOXC1
Formulation:	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Concentration:	lot specific
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Stability:	1 year
Predicted Protein Size:	Calculated MW: 57 kDa; Observed MW: 75 kDa
Gene Name:	forkhead box C1
Database Link:	Q12948



[View online »](#)


Background:

Swiss-Prot Acc.Q12948. DNA-binding transcriptional factor that plays a role in a broad range of cellular and developmental processes such as eye, bones, cardiovascular, kidney and skin development (PubMed:11782474, PubMed:15299087, PubMed:15684392, PubMed:16492674, PubMed:27907090, PubMed:14506133, PubMed:14578375, PubMed:15277473, PubMed:16449236, PubMed:17210863, PubMed:19793056, PubMed:19279310, PubMed:25786029, PubMed:27804176). Acts either as a transcriptional activator or repressor (PubMed:11782474). Binds to the consensus binding site 5'-[G/C][A/T]AAA[T/C]AA[A/C]-3' in promoter of target genes (PubMed:7957066, PubMed:11782474, PubMed:12533514, PubMed:14506133, PubMed:19793056, PubMed:27804176). Upon DNA-binding, promotes DNA bending (PubMed:7957066, PubMed:14506133). Acts as a transcriptional coactivator (PubMed:26565916). Stimulates Indian hedgehog (Ihh)-induced target gene expression mediated by the transcription factor GLI2, and hence regulates endochondral ossification. Acts also as a transcriptional coregulator by increasing DNA-binding capacity of GLI2 in breast cancer cells (PubMed:26565916). Regulates FOXO1 through binding to a conserved element, 5'-GTAAACAAA-3' in its promoter region, implicating FOXC1 as an important regulator of cell viability and resistance to oxidative stress in the eye (PubMed:17993506). Cooperates with transcription factor FOXC2 in regulating expression of genes that maintain podocyte integrity. Promotes cell growth inhibition by stopping the cell cycle in the G1 phase through TGFβ1-mediated signals (PubMed:12408963). Involved in epithelial-mesenchymal transition (EMT) induction by increasing cell proliferation, migration and invasion (PubMed:20406990, PubMed:22991501). Involved in chemokine CXCL12-induced endothelial cell migration through the control of CXCR4 expression. Plays a role in the gene regulatory network essential for epidermal keratinocyte terminal differentiation (PubMed:27907090). Essential developmental transcriptional factor required for mesoderm-derived tissues, such as the somites, skin, bone and cartilage. Positively regulates CXCL12 and stem cell factor expression in bone marrow mesenchymal progenitor cells, and hence plays a role in the development and maintenance of mesenchymal niches for haematopoietic stem and progenitor cells (HSPC). Plays a role in corneal transparency by preventing both blood vessel and lymphatic vessel growth during embryonic development in a VEGF-dependent manner. Involved in chemokine CXCL12-induced endothelial cell migration through the control of CXCR4 expression. May function as a tumor suppressor (PubMed:12408963).

Synonyms:

ARA; FKHL7; FREAC-3; FREAC3; IGDA; IHG1; IRID1; RIEG3

Product images:



Western blot detection of FOXC1 in 3T3 cell lysates using FOXC1 Rabbit mAb(1:1000 diluted). Predicted band size:57kDa. Observed band size:75kDa.